



ALMONDS: NUTRITION AND SCIENTIFIC RESEARCH

UPDATED SEPTEMBER 2017




ALMONDS: NUTRITION AND SCIENTIFIC RESEARCH

For more than two decades, Almond Board of California has invested in sound science to better understand the nutrient composition and health benefits of almonds. The ever-expanding body of almond nutrition research totals over 145 scientific publications to date, in areas including heart health, weight management, diabetes, nutrient composition and diet quality. Growing interest in weight management and smart snacking has prompted a shift in emphasis from the well-established body of evidence on heart health toward diabetes, weight management, satiety and cognitive

research to support a healthy lifestyle. (See Figure 1.)

The powerful nutrient package of almonds—low on the glycemic index and providing 6 grams of plant based protein, 4 grams of filling dietary fiber, 13 grams of unsaturated fats, and important vitamins and minerals including vitamin E (50% DV), magnesium (20% DV), and potassium (4% DV) per one-ounce handful, about 23 almonds—makes them an ideal fit for healthy lifestyles and a deliciously easy way to snack smarter.

Figure 1: Concerns about health condition
(n = 5,500 global consumers)



Source: Almond Board Global Perceptions Study, 2013.
Question: For the following health concerns, please indicate how impacted you are by each. (select all that apply.)

ALMONDS: A HEART-SMART SOLUTION

OVER TWO DECADES OF RESEARCH SHOWS THAT ALMONDS CAN HELP MAINTAIN A HEALTHY HEART AND HEALTHY CHOLESTEROL LEVELS.

Although heart disease remains the number one cause of death in the United States and worldwide, it is estimated that at least 80% of premature deaths from cardiovascular disease can be avoided if diet and lifestyle risk factors are controlled.¹ Diet is integral to managing cardiovascular risk, and more than two decades of research support the role of almonds in helping to maintain a healthy heart. In fact, the U.S. Food and Drug Administration says that scientific evidence suggests, but does not prove, that eating 1.5 ounces of almonds as part of a diet low in saturated fat and cholesterol may reduce the risk of heart disease.

Many randomized controlled studies have been conducted to examine the impact of almond consumption on markers of heart health, such as total and LDL cholesterol, HDL cholesterol, abdominal fat, oxidative stress and inflammation.

ALMONDS AND CHOLESTEROL

A 2016 meta-analysis and systematic review examined the breadth of research on almonds and heart health. The analysis of 18 published randomized controlled trials with a total of 837 participants showed significant favorable effects of almonds

PROTEIN
6g

VITAMIN E
7.3mg
50% DV

FIBER
4g
13% DV

UNSATURATED FATS
13g

POTASSIUM
210mg
4% DV

MAGNESIUM
76mg
20% DV



1 OZ = 23 ALMONDS



on total cholesterol, LDL cholesterol and triglycerides, with no change in HDL cholesterol levels? The effects of almonds on total cholesterol were dose-dependent, with a larger almond intake resulting in a greater reduction in total cholesterol. The evidence strongly indicates that almonds should be encouraged as part of a healthy diet to help maintain healthy blood lipid levels and reduce the risk of heart disease.

HDL CHOLESTEROL AND ALMONDS

In general, cholesterol-lowering diets reduce HDL cholesterol. However, studies show that when almonds are included in such diets, there is no significant effect—protective HDL is preserved?

Research supporting the role of almonds in heart health began in 1992, with the first study demonstrating that an almond-based diet (with 100 grams or 3.5 ounces of almonds per day) improved cholesterol levels.³ This landmark study helped set the stage for the almond nutrition research

program and provided compelling evidence that despite their high fat and calorie content, almonds could be included as part of a heart-healthy diet.

In the majority of studies of people with high cholesterol, the daily consumption of one to four ounces (28 to 114 grams) of almonds per day resulted in significant reductions in total and LDL cholesterol levels.^{4,5,6} In one study, 27 adults with high cholesterol ate heart-healthy diets with one of three snacks over a three-month period: 2.8 ounces (79 grams) of almonds, 1.3 ounces (37 grams) of almonds or a low-saturated-fat whole-wheat muffin as a daily snack⁶ (See Figure 2.) Researchers found that participants lowered their LDL cholesterol level an average of 4.4% with the 1.3-ounce (37 gram) portion of almonds and 9.4% with the 2.8-ounce (79 gram) portion. These results suggest there is a “dose effect of almonds on cholesterol levels—that higher intakes are associated with greater cholesterol-lowering effects.”

Another year-long crossover study on 81 U.S. adults (43 men and 38 women, age 49 years,

BMI 25 kg/m²) found that daily consumption of usual diet plus 52 grams (1.8 ounces) per day of almonds for six months without any additional dietary advice vs. their usual diet without almonds improved both TC:HDL and LDL:HDL ratios. Participants with high cholesterol levels showed greater response in changes in total cholesterol, LDL and TC:HDL and LDL:HDL ratios than those with normal cholesterol levels.⁷

Additionally, a four-week randomized analysis of previously collected data from 27 adults with elevated LDL showed that eating almonds daily as part of a healthy diet improved participants’ serum fatty acid profiles and reduced estimated (based on the Framingham equation) 10-year coronary heart disease risk scores by 3.5%.⁸ Limitations included lack of randomization in the diet order and lack of control over external factors that may have affected dietary behaviors over the course of the study⁷ and a relatively high dropout rate and potential confounding of MUFA intake.⁸

Two recent studies have examined the heart-health impact of swapping out higher carbohydrate snacks for almonds. In one study among Korean adults, researchers compared the effects of almond consumption versus cookies of equal caloric value as a daily snack on cardiovascular risk factors in 84 overweight and obese individuals.⁹ They found that participants in the almond group experienced a significant decrease in total cholesterol, LDL cholesterol and non-HDL cholesterol compared to the cookie group. Almonds also enhanced vitamin E status and serum total and LDL cholesterol in the overweight and obese individuals. Thus, including almonds as a snack can help healthy overweight/obese individuals improve nutritional status and reduce risk for CVD. This was the first study to look at almond consumption in a Korean population. Although most of the research has been conducted in North American and European populations, we have seen similar results from studies in Taiwan, India,

and now Korea, indicating that the heart-health benefits are similar between these genetically diverse groups.


In another study, 48 middle-aged women and men, normal and overweight, with elevated LDL cholesterol and normal HDL cholesterol levels at baseline, were assigned a cholesterol-lowering diet that either included almonds (43 grams or 1.5 ounces per day) or a calorie-matched high-carbohydrate snack (muffin).¹⁰ Researchers assessed almond consumption vs. high-carbohydrate snack on HDL particle type, distribution and transport of cholesterol to the liver for elimination. Compared with the control diet, the almond diet increased alpha-1 HDL (larger, more mature HDL; generally a marker of heart-health protection) as well as cholesterol efflux (removal of cholesterol from peripheral tissues for elimination), an important part of the cardio-protective role of HDL, in participants.

Research has also investigated the effects of almonds as part of a group of cholesterol-lowering foods, including plant sterols and soluble fiber. This diet, known as the Portfolio Eating Plan, consists of a National Cholesterol Education Program (NCEP) “Step 2” diet (saturated fat less than 7% of calories, less than 200 mg cholesterol) plus almonds (30 grams or 1 ounce per day); viscous fiber (20 grams per day), such as oats, barley, psyllium, legumes, eggplant and okra; vegetable protein (80 grams per day, half from soy), such as soy foods, beans, chick peas and lentils; and plant sterols (2 grams per day), such as plant sterol margarine. Initial studies that were done in a controlled environment (with all meals provided) resulted in LDL cholesterol reductions of nearly 30% in study participants, 46 adults with high cholesterol.¹¹ Later studies conducted in people with high cholesterol who followed the Portfolio Eating Plan on their own showed more modest,

but still significant, reductions in LDL, with an average reduction of 13% after one year of following the diet. The results from this collection of studies demonstrate that almonds can be part of a heart-healthy diet that—in combination with other cholesterol-improving components—effectively lowers cholesterol in study participants with high cholesterol.

While the majority of studies have been conducted in people with elevated cholesterol levels, there have also been studies investigating the impact of eating almonds on cholesterol in healthy people. The studies suggest that almonds have no detrimental effects on blood lipids and in fact, in one controlled-feeding study, eating 2.4 ounces (68 grams) of almonds per day for a period of four weeks actually improved the blood lipid profiles of healthy men and women by significantly reducing total and LDL cholesterol levels and improving

Figure 2: Change from baseline at four weeks in blood lipids on control, half-dose almonds and full-dose almonds. Values are mean ± SEM





ALMONDS AND DIABETES

THE UNIQUE NUTRIENT PACKAGE IN ALMONDS MAKES THEM A SMART CHOICE FOR MANAGING HEALTHY BLOOD SUGAR LEVELS.

The prevalence of type 2 diabetes is rapidly increasing. According to the most recent 2014 report from the Centers for Disease Control and Prevention, more than 29 million Americans have diabetes, up from 26 million in 2010. One in four people have diabetes and don't know it. And another 86 million adults—more than one in three U.S. adults—have prediabetes.¹⁷ Diabetes is also a contributing risk factor for other chronic diseases, such as heart disease and stroke. Dietary and lifestyle interventions are a critical component of diabetes management, and evidence continues to mount supporting the role of almonds and other tree nuts as part of an overall dietary pattern that is beneficial for those with type 2 diabetes. The nutrient profile of almonds—low-glycemic index and providing a satisfying combination of protein (6 grams per ounce), fiber (4 grams per ounce) and monounsaturated fats—makes them an ideal snack and addition to meals for individuals with impaired glucose intolerance or type 2 diabetes.

Many randomized controlled studies have been conducted to examine eating almonds in relation to blood glucose control. These studies were conducted in different population groups, including people with normal blood glucose control, people with prediabetes and people with type 2 diabetes (T2D).

IMPACT OF ALMONDS IN PARTICIPANTS WITH TYPE 2 DIABETES

A number of randomized, controlled studies of the effects of almonds on measures related to blood glucose control have been conducted in participants with T2D, evaluating both post-meal effects and longer-term measures (over at least four weeks). In four of the five longer-term studies, eating an almond-enriched diet resulted in significant reductions in fasting glucose and insulin levels control, when compared to an almond-free diet. One randomized trial in 19 U.S. adults (including 7 with T2D) reported a 30% reduction in postprandial glycemia in participants with T2D following the consumption of a test meal containing one ounce (28 grams) of almonds compared to an almond-free test meal similar in calories, fat and available carbohydrate, although the effect was not significant in those without T2D.¹⁸ These




the ratio of LDL to HDL compared to no almond consumption, a remarkable finding given that all participants were consuming a low saturated fat, National Cholesterol Education Program (NCEP) Step I diet.¹² A second study assessed the effect of eating a low-almond diet (10% of calories), a high-almond diet (20% of calories) or control diet (no almonds) in 16 healthy men and women (mean age 41 years).¹³ The high-almond diets significantly lowered average total cholesterol (-10 mg/dL) and LDL cholesterol (-10 mg/dL) compared to the control diets, while also increasing vitamin E levels in a dose-response manner.

EMERGING RISK FACTORS: INFLAMMATION AND BELLY FAT

Several studies have investigated the effects on emerging risk factors for cardiovascular disease such as inflammation and abdominal (belly) fat. One randomized controlled crossover feeding study assessed the effects of almonds on markers of inflammation in 25 healthy adults (ages 22 to 53). Participants were fed three different diets for four weeks each: a heart-healthy control diet (no nuts, <30% of calories from fat), a moderate-almond diet (10% of calories from almonds) and a high-almond diet (20% of calories from almonds).¹⁴ E-selectin (an inflammation marker) decreased as percentage of energy from almonds increased. C-reactive protein (another inflammatory marker) was lower in both almond diets compared to the control diet, and E-selectin decreased as percentage of energy from almonds increased. While not all markers of inflammation were improved, these findings suggest including almonds in a heart-healthy diet may help improve two important markers, C-reactive protein and E-selectin, and in turn, contribute to the prevention of heart disease.

In another study, 30 normal-weight Iranian men with mildly elevated cholesterol consumed 60 grams (2 ounces) of almonds daily for four weeks in addition to their usual diets. After four weeks, eating

Figure 3: Effects of eating 1.5 ounces of almonds per day compared to 150-calorie muffin on lipid profiles in adults with high cholesterol from baseline to six weeks



almonds significantly decreased total and LDL cholesterol, as well as apolipoprotein B100, a protein that plays a role in moving cholesterol throughout the body and a form of LDL cholesterol.¹⁵ ApoB 100 is thought to be an important determinant of cardiovascular risk. Almond consumption was also associated with an improvement in lipid oxidation parameters, suggesting that almonds may reduce the ability of fats to become oxidized in the body, a process that can lead to increased heart disease risk.

A study from Penn State University showed that snacking on almonds daily for six weeks not only reduced LDL and total cholesterol but also reduced abdominal fat and waist circumference in study participants.¹⁶ During this study, 52 adult participants (who were overweight with elevated LDL and total cholesterol but otherwise healthy) ate standard healthy diets that were identical except for the snack, either 1.5 ounces (43 grams) of almonds or a high-carbohydrate muffin with the same number of calories. Compared with snacking on muffins, eating

almonds significantly decreased total cholesterol (-5.1 mg/dL) and LDL cholesterol (-5.3 mg/dL) and maintained HDL cholesterol. HDL cholesterol actually decreased among participants in the muffin group. (See Figure 3.) There was a small weight loss in both groups that did not differ between the diets, but snacking on almonds actually reduced abdominal fat (-0.15 pounds) and waist circumference (-0.31 inches) compared with snacking on muffins. The overall diets were not matched for macronutrient content. The study suggests that regularly choosing almonds instead of a high-carb snack may be a simple dietary strategy to help improve body composition.

CONCLUSIONS

Dietary changes are often the first and one of the most effective ways to reduce the risk of cardiovascular disease, and the body of research suggests that eating almonds can help maintain a healthy heart and healthy cholesterol levels.

same researchers conducted a pilot study on the longer-term effects of almonds on glucose control in 13 adults with T2D. Participants consumed a daily one-ounce serving of almonds (five days per week for 12 weeks) or a cheese snack with the same number of calories. After 12 weeks, hemoglobin A1c in individuals with T2D was reduced by 4% in participants who consumed almonds daily compared to baseline.

Another 10-week crossover trial in 20 Chinese adults with T2D and mild hyperlipidemia (9 male, 11 female; age 58 years; BMI 26kg/m²) investigated the effects of a four-week diet containing two ounces (56 grams) of almonds per day versus a no-almond control diet.¹⁹ The study demonstrated that almond consumption

helped improve glycemic control by lowering fasting insulin and fasting glucose as well as decreasing the risk for heart disease through significant reductions on total cholesterol (-6%), LDL cholesterol (-11.6%) and LDL:HDL ratio when compared to the control. In a third long-term study, participants (65 overweight and obese adults) consumed three ounces (85 grams) of almonds per day as part of a healthy diet for 12 weeks and showed improvements in HbA1c.²⁰

Finally, a recent six-month study among 50 Asian Indians with T2D and elevated cholesterol examined the effect of almonds on CVD risk factors.²¹ During a three-week run-in period, participants ate a standard diet compliant with the dietary guidelines for Asian Indians and appropriate for diabetes. During this period, participants were also



asked to walk for 45 minutes at least five days a week to standardize their physical activity and were instructed to maintain the same level of activity for the rest of the study. Whole natural unroasted almonds (20% of energy intake) were substituted for fat (such as cooking oil and butter) and some carbohydrates in the intervention group. Following the almond intervention, waist circumference, waist-to-height ratio, total cholesterol, triglycerides, LDL cholesterol, C-reactive protein (an indicator of inflammation) and hemoglobin A1C (an indicator of long-term blood sugar control) improved in participants. The study findings illustrate that incorporation of almonds in a well-balanced healthy diet leads to multiple beneficial effects on glycemic and cardiovascular risk factors.

The summary of these results suggest that modest almond consumption improves both short-term and long-term markers of glucose control in individuals with T2D. The studies were well controlled and of sufficient duration to determine effects on glycemic control; they are limited by their small sample size and, in some cases, have limited generalizability to free-living conditions in the studies in which meals were provided to participants.

ALMONDS AND PREDIABETES

Studies also suggest that almonds may have benefits for people with prediabetes. One short-term post-meal study in 14 adults with impaired glucose tolerance showed that the consumption of a 580-kcal breakfast meal containing 1.5 ounces (44.5 grams) of almonds resulted in significant reductions in study participants' blood glucose levels both acutely after breakfast and after a second meal relative to the consumption of a 347-kcal control breakfast meal, which differed in total dietary energy but provided the same amount of available carbohydrate.²² A long-term 16-week randomized control trial on 65 middle-aged U.S. adults (48 women and 17 men) with prediabetes investigated the effects of consuming an American

Diabetes Association diet consisting of 20% of calories from almonds (approximately 2 ounces (57 grams) per day) on the progression of T2D and CVD. The group that consumed the almond-enriched diet showed significantly improved LDL cholesterol levels and measures of insulin sensitivity, both of which are risk factors for heart disease and T2D.²³ (See Figure 4.) The study was of sufficient duration to examine effects on markers of long-term blood glucose control; however, reliance on a single fasting sample for measurement of insulin resistance is an analytical limitation.


EFFECT OF ALMONDS ON BLOOD GLUCOSE MEASURES IN HEALTHY PEOPLE

Post-meal studies conducted in healthy or hyperlipidemic participants with normal blood glucose control cumulatively suggest that almonds have neutral or beneficial effects on post-meal blood glucose and insulin responses; in some studies, almonds actually reduced post-meal blood glucose and insulin spikes as well as blood glucose and insulin levels over a two-hour time period relative to an almond-free meal.^{24,25}

In longer-term four-week studies, eating about 1.2 or 2.5 ounces (35 or 70 grams) of almonds per day resulted in significant reductions in a marker of insulin secretion, suggesting a decrease in insulin resistance²⁶ as well as significant dose-dependent improvements in total cholesterol and other blood lipids.²⁷ Calorie intake was similar between the control and almond diets in both studies, but the duration was too short to discern effects on long-term blood glucose control.

CONCLUSIONS

Based on the totality of scientific evidence from randomized controlled studies, almonds, when eaten as part of a healthy diet, may have beneficial effects on blood glucose and insulin responses, both in the short term after consuming a meal and over the longer term, especially in those with impaired glucose tolerance and/or T2D. Dietary changes are often the first and one of the most effective ways to manage diabetes, and the body of research suggests that eating almonds can help maintain healthy blood sugar levels.



ALMONDS: A SATISFYING WEIGHT-WISE SNACK

A DAILY HANDFUL OF ALMONDS IS A DELICIOUS WAY TO MANAGE CRAVINGS AND HELP MAINTAIN A HEALTHY BODY WEIGHT.

The prevalence of overweight and obesity continues to be a major public health issue worldwide. According to data from the National Health and Nutrition Examination Survey (NHANES), 2013–2014, in the U.S., 70% of men and women—more than two in three adults—are considered to be overweight or have obesity and one in six children and adolescents (17%) are obese.²⁸ Given that snacking has become nearly universal behavior, with an estimated 94% of Americans reporting snacking at least once a day²⁹ combined with persistently high obesity rates, identifying nutrient-rich snack options that pose little risk for weight gain is of growing importance. The nutrients in almonds, including monounsaturated fat (13 grams per ounce) and fiber (4 grams per ounce), are associated with improved satiety, suggesting they would be an ideal snack for those concerned about weight management.

New data from the U.S. Department of Agriculture (USDA) shows that both roasted and unroasted almonds provide fewer calories than thought—and that the number of calories is largely dependent on form.³⁰ (See Figure 5.) The study, conducted by scientists from USDA's Agricultural Research Service (ARS) and jointly funded by Almond Board of California and USDA ARS, shows that compared to the number of calories listed on nutrition labels, participants actually absorbed 25% fewer calories from whole unroasted almonds, 19% fewer calories from whole roasted almonds and 17% fewer calories when almonds were roasted and chopped. Measured calories in almond butter did not differ from calories listed on labels. Much of this finding has to do with particle size after chewing and digestion. The larger the particle size, after chewing for example, the less the almond is able to be broken down by digestive enzymes and more of the almond is excreted, so fewer calories are absorbed. The reverse is also true: the smaller the particle size, the more almond cells are exposed to digestive enzymes and the more calories are absorbed. In addition to chewing and digestion, mechanical processes, such

as chopping, grinding and roasting almonds can also impact particle size.


Further research is needed to better understand how this technique for calculating calories could potentially affect the calorie content of other foods.

Many randomized controlled studies have been conducted to examine the effects of almonds, consumed as part of a sensible eating plan, on outcomes related to satiety (like hunger, fullness, desire to eat and prospective food consumption) and/or body composition (like body weight, body mass index (BMI), body fat and waist circumference). These studies were conducted in different population groups, including people with normal weight, as well as overweight or obese people.

IMPACT OF EATING ALMONDS ON MEASURES OF HUNGER, SATIETY AND SUBSEQUENT CALORIE INTAKE IN NORMAL-WEIGHT PEOPLE

In post-meal studies, the daily consumption of almonds is associated with improving ratings of hunger and satiety in healthy people. In one study, the daily consumption of 2.8 ounces (80.4 grams) of almonds reduced subjective rating of hunger³¹ and a second study investigated the effects of two different portion sizes of almonds (1 and 1.5 ounces or 28 and 43 grams) as a midmorning snack on satiety and energy intake, in comparison to having no snack. (See Figure 6.) There were no significant differences in total daily energy intake between any of the groups, indicating that participants—healthy, Caucasian women—naturally compensated for the almond calories consumed, whether they had one (160 calories) or 1.5 servings (250 calories) of almonds as the mid-morning snack.³² After eating their usual breakfast and having the mid-morning almond snack, participants were fed lunch midday and permitted to eat as much as they wanted until they were comfortably full. Ratings of appetite and fullness were dose-dependent,


Figure 5: Food processing and structure impact the metabolizing energy of almonds.



with participants reporting being the least hungry when they ate 1.5 ounces (43 grams) of almonds and the hungriest on the day when they didn't eat almonds. Although habitual almond intake was not controlled for and a control snack was not tested, the studies suggest that snacking on nutrient-rich almonds may improve satiety and help control cravings.

A longer-term four-week randomized, controlled clinical study in which 137 adults who were at risk of T2D but otherwise healthy were given 1.5 ounces (43 grams) of almonds as a snack or with meals also showed significantly greater reductions in daylong ratings of hunger and desire to eat in participants who consumed almonds either a snack or part of a meal relative to those who did not consume almonds.³³

Figure 6: A midmorning snack of almonds generates satiety





Despite consuming approximately 250 calories from almonds every day for over four weeks, participants did not increase their daily total calorie intake or experience any change in weight over the course of the study. Although the study was of relatively short duration, these findings suggest almonds may be a satisfying snack option to help maintain a healthy weight.

In another 10-week study, 20 healthy adult women consumed their normal diets plus 344 calories (approximately 2 ounces or 56 grams) of almonds per day for 10 weeks and then followed their normal diet without almonds for 10 weeks, with a three-week washout period in between (See Figure 7).³⁴ There were no differences in body weight, metabolic rate or energy expenditure observed, suggesting that the almonds replaced other foods in the diet and, therefore, did not increase overall calorie intake.

EFFECTS OF EATING ALMONDS ON SATIETY AND WEIGHT IN PEOPLE WHO ARE OVERWEIGHT OR OBESE

There have been a number of studies investigating the short-and long-term effects of almonds on measures related to body composition and weight in overweight and obese adults (BMI \geq 25 kg/m²). In one study, overweight women who consumed a meal containing 1 ounce (28 grams) of almonds reported feeling more hungry, less full, more desire to eat, with a greater food consumption later in the day relative to when they consumed a control meal containing a mixture of safflower and corn oils, although differences in satiety ratings were not observed in overweight men.³⁵ In another study conducted in obese adults, the consumption of a meal containing 1.5 ounces (43 grams) of almonds boosted fullness ratings in the afternoon and throughout the day relative to the consumption of a control meal without almonds, which was lower in total dietary energy but provided the same amount of available carbohydrate.³⁶

Long-term studies cumulatively suggest that almonds have no detrimental effects on body composition in overweight or obese participants; in fact, significant improvements in body composition have been observed in two studies.

In obese adults with type 2 diabetes, the consumption of one ounce (28 grams) of almonds five days per week for 12 weeks resulted in a significant decrease in BMI relative to no almond consumption.³⁷ In another study, 65 overweight and obese participants who consumed three ounces (84 grams) of almonds daily for 24 weeks had significantly greater reductions in body weight, BMI, waist circumference, fat mass and total body water compared to participants who did not consume almonds.³⁸ (See Figure 8.) These findings are remarkable, given that participants consumed the almonds as part of a healthy low-calorie diet.

Another 18-month clinical trial examined the effects of a low-calorie diet containing two ounces (56 grams) per day of almonds compared to a low calorie, nut-free diet on weight loss and heart disease risk

factors in 123 overweight or obese adults. Although both groups had lost similar and significant amounts of weight after 18 months, compared with the nut-free group, the almond-enriched diet was associated with greater reductions in total cholesterol, TL:HDL ratio, and triglycerides.³⁹ Strengths of the long-term studies include sufficient study duration to discern the effects on body weight and adequate control of the total energy intake between the control and almond diets. The studies are limited by inadequate control of the confounding effects of physical activity.

CONCLUSIONS

The body of scientific evidence suggests that despite their relatively high energy density, almonds, when eaten as a part of a healthy diet, do not cause weight gain and may even have beneficial effects on body composition, especially in overweight or obese adults. Several mechanisms give explanation for the positive associations between almonds and other nuts and energy balance and body weight, including their strong satiety effects, incomplete calorie availability and possible enhancement of resting energy expenditure.⁴⁰ Although many commonly consumed

snacks provide empty calories, almonds are a healthy, nutrient rich snack choice. The unique nutrient package in almonds makes them a satisfying, weight-wise snack.

ALMONDS AND DIET QUALITY

The 2015 Dietary Guidelines for Americans recommend improving diet quality; and one way to do this is by replacing high-calorie snacks with nutrient-rich options. A recent study that evaluated the potential effects of replacing typical snack foods with almonds and other tree nuts shows that this simple swap would decrease empty calories, solid fats, saturated fat and sodium in the diet, while increasing intake of key nutrients.⁴¹ Using data of more than 17,000 children and adults from the National Health and Nutrition Examination Survey (NHANES; 2009–2012), the researchers applied food pattern modeling to assess the hypothetical impact of replacing all snack foods, excluding beverages, with tree nuts (model 1) and replacing all but “healthy” snack foods (whole grains, whole fruits and non-starchy vegetables) with tree nuts (model 2). Almonds are the most frequently consumed nut and in this study, 44% of all tree nuts eaten were almonds. Therefore, assessments using the NHANES data were repeated using almonds only. All reported snacks were replaced calorie for calorie with almonds or other tree nuts, reflecting typical American consumption patterns. The Healthy Eating Index 2010, which measured adherence to



the 2010 Dietary Guidelines for Americans, was used to assess diet quality.

In both models examined, where tree nuts hypothetically replaced all snack foods and where tree nuts hypothetically replaced only less-healthy snack foods, consumption of empty calories, solid fats, saturated fat, sodium, carbohydrates and added sugars all declined, while consumption of oils and good fats increased significantly. Fiber and magnesium also increased, while protein increased by a small margin. The findings were the same in the almond-only model.

This study echoes findings from a similar NHANES analysis on almond eaters, which examined the characteristics of almond eaters. It found that people who reported eating almonds had higher intake of key nutrients (such as dietary fiber, calcium, potassium and iron, as well as higher intakes of several other “shortfall nutrients,” including vitamins A, D, E and C; folate; and magnesium), better overall diet quality (measured by Healthy Eating Index scores) and lower body mass index and waist circumference compared to non-consumers.⁴² Almond consumers (defined as those eating about 1 ounce (28 grams) per day) also tended to be more physically active and less likely to smoke than their non-almond-eating counterparts, suggesting that including almonds as a regular part of the diet is associated with a portfolio of healthy lifestyle attributes.

Another study on diet quality conducted by the University of Florida examined the impact of eating almonds and/or almond butter for three weeks on diet quality in addition to microbiota composition in 29 parents and their children. Participants ate either 1.5 ounces (43 grams) of almonds and/or 0.5 ounces (14 grams) of almond butter on a daily basis for three weeks, as part of their usual diet. This was followed by a four-week washout period and a three-week control period in which no almonds were eaten.

Diet quality was assessed based on the U.S. Dietary Guidelines. When parents and children ate almonds, their overall diet quality improved, as measured by an increase in Healthy Eating Index (HEI) scores (a standard measure of adherence to recommended dietary guidance).⁴³ Specifically, parent and child scores increased for fatty acids, total protein, seafood and plant protein, and decreased for fruit and empty calories. In addition, when eating almonds, participants also consumed significantly more vitamin E and magnesium, two nutrients commonly underconsumed by the majority of adults and children. Although no specific changes in immune markers were observed, almond consumption did result in detectable changes in gut microbiota. More research is necessary to understand these changes and their potential health impacts.



- World Heart Federation, "About World Heart Day," Web. 24 September 2014. <http://www.world-heart-federation.org/what-we-do/awareness/world-heart-day-2014-home/about-world-heart-day/>.
- Musa-Veloso K, Paulionis L, Poon T, Lee HL. The effects of almond consumption on fasting blood lipid levels: a systematic review and meta-analysis of randomised controlled trials. *Journal of Nutritional Science* 2016; 5(e34): 1-154.
- Spiller, GA et al. Effect of a diet high in monounsaturated fat from almonds on plasma cholesterol and lipoproteins. *Journal of the American College of Nutrition*. 1992 Apr;11(2): 126-30.
- Spiller GA, Jenkins DA, Bosello O, Gates JE, Cragen LN, Bruce B. Nuts and plasma lipids: an almonds-based diet lowers LDL-C while preserving HDL-C. *Journal of the American College of Nutrition*. 1998; 17(3): 285-90.
- Tamizfar et al., 2005.
- Jenkins DJ, Kendall CW, Marchie A, Parker TL, Connelly PW, Qian W, Haight JS, Faulkner D, Vidgen E, Lapsley KG, Spiller GA. Dose response of almonds on coronary heart disease risk factors: blood lipids, oxidized low-density lipoproteins, lipoprotein(a), homocysteine, and pulmonary nitric oxide: a randomized, controlled, crossover trial. *Circulation*. 2002; 106(11): 1327-32.
- Jaceldo-Siegl K, et al. Influence of body mass index and serum lipids on the cholesterol-lowering effects of almonds in free-living individuals. *Nutrition, Metabolism and Cardiovascular Diseases*. 2011; 21: S7-S13.
- Nishi S, Kendall CW, Gascoyne AM, et al. Effect of almond consumption on the serum fatty acid profile: a dose response study. *British Journal of Nutrition* 2014, 1-10. doi:10.1017/S0007114514001640.
- Jung H, Chen C-Y, Blumberg JB, Kwak HK. The effect of almonds on vitamin E status and cardiovascular risk factors in Korean adults: a randomized clinical trial. *European Journal of Nutrition*. 2017. doi: 10.1007/s00394-017-1480-5.
- Berryman CE, Fleming JA, Kris-Etherton PM. Inclusion of almonds in a cholesterol-lowering diet improves plasma HDL subspecies and cholesterol efflux to serum in normal-weight individuals with elevated LDL cholesterol. *The Journal of Nutrition* 2017; doi: 10.3945/jn.116.245126.
- Jenkins DJ, Kendall CW, Marchie A, Faulkner DA, Wong JM, de Souza R, Emam A, Parker TL, Vidgen E, Lapsley KG, Trautwein EA, Josse RG, Leiter LA, Connelly PW. Effects of a dietary portfolio of cholesterol-lowering foods vs lovastatin on serum lipids and C-reactive protein. *Journal of the American Medical Association*. 2003 July 23; 290(4): 502-10.
- Sabaté J, Haddad E., Tanzman JS., Jambazian P., Rajaram S. (2003). Serum lipid response to the graduated enrichment of a Step I diet with almonds: a randomized feeding trial. *American Journal of Clinical Nutrition*; 77(6): 1379-1384.
- Jambazian PR, Haddad E, Rajaram S, Tanzman J, Sabaté J. Almonds in the diet simultaneously improve plasma alpha-tocopherol concentrations and reduce plasma lipids. *Journal of the American Dietetic Association* 2005; 105(3): 449-54.
- Rajaram S, Connell KM, Sabaté J. 2010. Effect of almond-enriched high monounsaturated fat diet on selected markers of inflammation: a randomized, controlled, crossover study *British Journal of Nutrition* 2010; 103: 907-912.
- Jalali-Khanabadi BA, Mozaffair-Khosravi H, Parsaeyan N. Effects of almond dietary supplementation on coronary heart disease lipid risk factors and serum lipid oxidation parameters in men with mild hyperlipidemia. *Journal of Alternative Complementary Medicine* 2010; 16(12): 1-5.
- Berryman CE, West SG, Fleming JA, Bordi PL, Kris-Etherton PM. Effects of Daily Almond Consumption on Cardiometabolic Risk and Abdominal Adiposity in Healthy Adults with Elevated LDL-Cholesterol: A Randomized Controlled Trial. *Journal of the American Heart Association* 2015; 4:e000993 DOI: 10.1161/JAHA.114.000993
- Diabetes Latest, Centers for Disease Control and Prevention. June 17, 2014. <https://www.cdc.gov/features/diabetesfactsheet>.
- Cohen A, et al. Almond ingestion at mealtime reduces postprandial glycemia and chronic ingestion reduces hemoglobin A1c in individuals with well-controlled type 2 diabetes mellitus. *Metabolism*. 2011; 60(9): 1312-1317.
- Li S, et al. Almond consumption improved glycemic control and lipid profiles in patients with type 2 diabetes mellitus. *Metabolism*. 2011; 60(4): 474-479.
- Wien MA, Sabate JM, Ikle DN, Cole SE, Kandeel, FR. (2003). Almonds vs. complex carbohydrates in a weight reduction program. *Int J. Obes. Relat. Metab. Disord*. 27(11): 1365-1372.
- Gulati S, Misra A, Pandey RM. Effect of almond supplementation on glycemia and cardiovascular risk factors in Asian Indians in North India with type 2 diabetes mellitus: A 24-week study. *Journal of Metabolic Syndrome and Related Disorders*. Epub ahead of print. Jan 4 2017. doi: 10.1089/met.2016.0066.
- Mori A, et al. Acute and second-meal effects of almond form in impaired glucose tolerant adults: a randomized crossover trial. *Nutr Metab (Lond)*. 2011; 8(1):6.
- Wien M, et al. Almond consumption and cardiovascular risk factors in adults with prediabetes. *Journal of the American College of Nutrition*. 2010; 29(3): 189-197.
- Josse AR, Kendall CWC, Augustin LSA, Ellis PR, Jenkins DJA. (2007). Almonds and postprandial glycemia—a dose-response study. *Metabolism* 56 (3): 400-404.
- Jenkins DJA, Kendall CWC, Josse AR, Salvatore S, Brighenti F, Augustin LSA, Ellis PR, Vidgen E, Rao AV. (2006). Almonds decrease postprandial glycemia, insulinemia, and oxidative damage in healthy individuals. *J. Nutr*. 136 (12):2987-2992.
- Jenkins DJA, Kendall CWC, Marchie A, Josse AR, Nguyen TH, Faulkner DA, Lapsley KG, Singer W. (2008). Effect of almonds on insulin secretion and insulin resistance in nondiabetic hyperlipidemic subjects: a randomized controlled crossover trial. *Metabolism* 57 (7): 882-887.
- Sabaté J, Haddad E, Tanzman JS, Jambazian P, Rajaram S. (2003). Serum lipid response to the graduated enrichment of a Step I diet with almonds: a randomized feeding trial. *American Journal of Clinical Nutrition*. 77 (6): 1379-1384.
- National Institute of Diabetes and Digestive and Kidney Diseases Overweight & Obesity Statistics. <https://www.niddk.nih.gov/health-information/health-statistics/overweight-obesity>. Accessed September 5, 2017.
- Mintel. Snacking Motivations and Attitudes U.S. April 2015. <http://store.mintel.com/snacking-motivations-and-attitudes-us-april-2015> Accessed September 5, 2017.
- Gebauer SK, Novotny JA, Bornhorst GM, Baer DJ. Food processing and structure impact the metabolizable energy of almonds. *Food & Function*. Published online 28 September 2016. DOI 10.1039/c6fo01076h.
- Kirkmeyer SV, Mattes RD (2000). Effects of food attributes on hunger and food intake. *Int J Obes Relat Metab Disord* 24(9): 1167.
- Hull S, Re R, Chambers L, Echaniz A, Wickham SJ. A midmorning snack generates satiety and appropriate adjustment of subsequent food intake in healthy women. *European Journal of Nutrition* 2014. DOI 10.1007/s00394-014-0759-2.
- Tan YT, Mattes RD. Appetitive, dietary and health effects of almonds consumed with meals or as snacks: a randomised, controlled trial. *European Journal of Clinical Nutrition* 2013; 67: 1205-14.
- Hollis J, Mattes R. Effect of chronic consumption of almonds on body weight in healthy humans. *Br J Nutr* 2007; 98: 651-656.
- Burton-Freeman B, Davis PA, Schneeman BO (2004). Interaction of fat availability and sex on postprandial satiety and cholecystokinin after mixed-food meals. *Am J Clin Nutr* 80: H1207-1214.
- Mori AM, Considine RV, Mattes RD (2011). Acute and second-meal effects of almond form in impaired glucose tolerant adults: a randomized crossover trial. *Nutr Metab (Lond)* 8(1):6. doi: 10.1186/1743-7075-8-6.
- Cohen AE, Johnston CS. Almond ingestion at mealtime reduces postprandial glycemia and chronic ingestion reduces hemoglobin A1c in individuals with well-controlled type 2 diabetes mellitus. *Metabolism* 2011; 60: 1312-1317.
- Wien MA, Sabate JM, Ikle DN, Cole SE, Kandeel FR. Almonds vs. complex carbohydrates in a weight reduction program. *Int J Obes Relat Metab Disord* 2003; 27(11): 1365-1372.
- Foster G, et al. A randomized trial of the effects of an almond-enriched, hypocaloric diet in the treatment of obesity. *American Journal of Clinical Nutrition*. 2012; 96(2): 249-254.
- Flores-Mateo G, et al. Nut intake and adiposity: meta-analysis of clinical trials. *Am J Clin Nutr*. 2013;97: 1346-55.
- Rehm CD, Drewnowski A. Replacing American snacks with tree nuts increases consumption of key nutrients among U.S. children and adults: results of an NHANES modeling study. *Nutrition Journal*. Published online March 7, 2017. DOI: 10.1186/s12937-017-0238-5.
- O'Neil CE, Nicklas TA, Fulgoni, III VL. Almond consumption is associated with better nutrient intake, nutrient adequacy and diet quality in adults: National Health and Nutrition Examination Survey 2001–2010. *Food and Nutrition Sciences* 2016.
- Burns AM, Zitt M, Rowe CC, Langkamp-Henken B, Volker M, Nieves Jr. C, Ukhanova M, Christman MC, Dahl WJ. Diet quality improves for parents and children when almonds are incorporated into their daily diet: a randomized, crossover study. *Nutrition Research* 2015; doi: 10.1016/j.nutres.2015.11.004.

